Dermatopathology Slide Review Part 113

Paul K. Shitabata, M.D.
Dermatopathology Institute
Torrance, CA











Syringoma arising with pigmented Seborrheic keratosis-Vulva


Seborrheic keratosis
Comma-Shaped ducts
Syringoma











What is the best diagnosis?

A. Mycosis fungoides
B. Lymphomatoid papulosis
C. Leukemia cutis
D. Anaplastic large cell lymphoma
E. Lichen planus

Mycosis fungoides

CD4



CD7



CD4 + with Loss of CD7 and CD5

Hyper-convolu-ted and hyper-chromatic lymphs



Pautrier Micro-abscess

Lymphs Intercalate with basal keratino-cytes












Giemsa Stain

What is the best diagnosis?

A. Amebiasis
B. Mucormycosis
C. Aspergillosis
D. Leishmaniasis
E. Verruga peruana

Leishmaniasis

Histiocytes with numerous Amasti-gotes



Suppura-tive necrosis













What is the best diagnosis?

A. Porocarcinoma
B. Nodular Hidradenoma
C. Proliferating Pilar Tumor
D. Nodular Basal Cell Carcinoma
E. Invasive Squamous Cell Carcinoma

Eccrine Porocarcinoma


Epidermal Attach-ment
Ductal Differen-tiation
Infiltrative Growth Pattern
Hyalinized Basement Membrane









What is the best diagnosis?

A. Pemphigus vulgaris
B. Epidermolysis bullosa acquisita
C. Dyshidrotic eczema
D. Bullous pemphigoid
E. Grover's disease

Dyshidrotic Eczema


Intraepidermal vesicles
Acral Skin
No acantholysis
Superficial perivascular
Lymphocytic infiltrate