

Disorders of Collagen

Weedon Chapter 11

Paul K. Shitabata, M.D.
Dermatopathology Institute

LSA vs. Morphea

Histopathology	LSA	Morphea
Elastic fibers decreased		
Interface dermatitis		
Thickened walls of small blood vessels		

Answers

Histopathology	LSA	Morphea
Elastic fibers decreased	+	
Interface dermatitis	+	
Thickened walls of small blood vessels		+

Correct associations with keloids include all except

- A. Lower rate of apoptosis
- B. Higher rate of p53 mutations
- C. Increased mast cells
- D. Dystrophic bone
- E. Histopathologic overlap with hypertrophic scars

Answer B

Lower rate of p53 mutations

All may show sclerodermoid-like plaques in the dermis except:

- A. Hutchinson-Gilford syndrome
- B. Atrophoderma of Pasini and Pierini
- C. Acrodermatitis Chronica Atrophicans
- D. Reactive perforating collagenosis

Answer D

- Reactive perforating collagenosis

Correct associations with chondrodermatitis nodularis helices include all except:

- A. Perichondrial fibrosis
- B. Eosinophilia of cartilage
- C. Ulceration and granulation tissue
- D. Atypical junctional melanocytic proliferation
- E. Association with autoimmune collagen vascular disease

Answer D

- Atypical junctional melanocytic proliferation