

Spirochetal Infections

(Weedon Chapter 24)

Paul K. Shitabata, M.D.

Dermatopathology Institute

Acceptable tissue stains to identify Spirochete organisms include all except:

- A. Barbour-Stoenner-Kelly
- B. Levaditi
- C. Warthin-Starry
- D. Antibodies to *T. pallidum*

Answer A

- A. Barbour-Stoenner-Kelly is a culture medium for Borrelia organisms.

Tissue histopathologic reactions associated with *Borrelia* infections include all except:

- A. Lymphoma
- B. Pseudolymphoma
- C. Dermal sclerosis
- D. Lichen sclerosus
- E. Acantholytic bullae

Answer E

- To date, no acantholytic bullous eruption has been associated with *Borrelia* infections.

Match the following:

| Disease | Histopathology |
|--------------------|--|
| Primary Syphilis | Caseating granulomas |
| Secondary Syphilis | Spirochetes predominately found in epidermis |
| Tertiary Syphilis | Psoriasiform hyperplasia |
| Yaws | Acute ulceration |

Match the following:

| Disease | Histopathology |
|--------------------|--|
| Primary Syphilis | Acute ulceration |
| Secondary Syphilis | Psoriasiform hyperplasia |
| Tertiary Syphilis | Caseating granulomas |
| Yaws | Spirochetes predominately found in epidermis |